



Cairo University Children Hospital
Abu El - Reech Hospital

Practical Guidelines
(Protocol)
for Management of Diarrhea
And Dehydration
In
Gastroenteritis Unit
Abu El - Reech Pediatric Hospital,
Al- Mounira
Faculty of Medicine
Cairo University

BY

Professor Dr Nabil Abdel Aziz

Director of Gastroenteritis Unit, Abu Elreech Hospital
Cairo University Children Hospital

Professor Dr Mona Abu Zekri

Former head of Gastroenteritis Unit, Abu Elreech Hospital
Cairo University Children Hospital

Dr Ahmed Youssef Ezz EIDin

Consultant Pediatrician
Gastroenteritis Unit, Abu Elreech Hospital
Cairo University Children Hospital

Dr Mazhar Ashour

Consultant Pediatrician
Gastroenteritis Unit, Abu Elreech Hospital
Cairo University Children Hospital

Dr Abeer Hussien

Pediatrician
Gastroenteritis Unit, Abu Elreech Hospital
Cairo University Children Hospital

Dr Mohamed Diab

Pediatrician
Gastroenteritis Unit, Abu Elreech Hospital
Cairo University Children Hospital

Acknowledgements

Dr. Ahmed Morad

Dr. Amr Mostafa

Dr. Ghada Fikry

Pediatricians G.E Unit, Abou El-Reech Hospital

جميع حقوق الطباعة محفوظة وتحت طائلة القانون

Diarrhea $\left\{ \begin{array}{l} \text{acute diarrhea} < 7 \text{ days} \\ \text{chronic diarrhea} > 15 \text{ days "continuous"} \end{array} \right.$

investig. & manage. \rightarrow تفحص وإدارة

Diarrhea is the major worldwide pediatric problem and it is serious problem in the developing countries.

Diarrhea is defined as: stools of increased frequency, fluidity and volume > 3 times/day. Or altered consistency of the stool.

How to deal with case of Diarrhea ^{Acute}

[I] Diagnosis:

1. Infectious:

- a) Viral. \rightarrow acute, low grade fever, no Tenosmus. *Stool analysis so H/O should be taken well.*
- b) Bacterial. *offensive, freq., grade fever, Mucous in blood.*
- c) Parasitic. *if D > 15 days bloody stool even for ID. patient < 5 Mo age old.*

2. Non-Infectious.

[II] Dehydration:

1. Degree:

- a) No dehydration.
- b) Some dehydration.
- c) Severe dehydration

2. Type (According to serum Na^+):

- a) Isotonic.
- b) Hypertonic.
- c) Hypotonic.

[III] Other Complications:

- Shock, Acidosis, Electrolytes disturbance, Bleeding, Hemolytic Uremic Syndrome (HUS), DIC and Convulsion.

Example for diagnosis:

Acute gastroenteritis, severe dehydration, shock, bleeding, DIC,

- Malnutrition "common" as $\left\{ \begin{array}{l} \downarrow \text{pta, vitamins, Minerals} \rightarrow \downarrow \text{villi} \\ \downarrow \text{immune defense} \rightarrow \text{susceptibility to infections} \end{array} \right.$
- Metabolic disorders e.g. lactose intolerance
 - Congenital: No lactase
 - Acquired: Malnutrition
 - 1- repeated infection \rightarrow destr. of brush border \rightarrow contains the lactase enzyme, \rightarrow give lactose free milk.

Nutrition
fluid replacement
Drugs?

Management of Diarrhea

- I - Rehydration + Feeding = Oral Rehydration Therapy (ORT).
- II - Treatment of infection.
- III - Management of complication.

To reach the proper diagnosis and management, perform the following:

I. History:

- Diarrhea, duration, frequency, consistency (\pm mucous, blood), *Tenesmus*
 - Vomiting, duration, frequency, character (i.e., projectile, regurge, coffee ground vomitus).
 - Fever.
 - Urinary flow.
 - Bleeding.
 - Convulsions.
 - **Feeding:**
 - 1) Breast milk.
 - 2) Formula, how many feds/day, how mixing with water.
 - 3) Weaning: type of foods administered.
 - **Medication:**
 - 1) Antibiotics.
 - 2) Antidiarrheal.
 - 3) ORS (amount – how to mix).
- mucous, blood, Tenesmus*
- MOT*
- water food Toilet*
- Guinea 2-3 yrs*
- Amebic dysentery*

II. Examination:

A. General Condition:

- Appearance.
- Breathing.
- Circulation to skin.

Ex.: Looks well, ill, sick, toxic, respiratory distress, cyanotic, pale, mottled skin.

B. Weight: for estimation of fluid intake and follow up.

C. Degree of dehydration:

1. No dehydration.

2. Some dehydration:

If the child has two or more of any of the following signs:

- Restlessness, irritable.
- Thirsty and drinks eagerly.
- Sunken eyes.
- Skin pinch goes back slowly.
- Depressed fontanel.

3. Severe dehydration:

Has two or more of the following signs:

- Lethargic.
- Unconsciousness.
- Sunken eyes.
- Drinks poorly, or unable to drink.
- Capillary refill (>2 sec).
- Cold extremities.
- Skin pinch goes back very slowly.

III. Systemic examination:

1. **Chest-heart:**


For chest infection, abnormal sounds, tachycardia, murmurs.

2. **Abdomen:**

For masses, distention, organomegaly.

3. **CNS:**

Convulsions (tonic, clonic, tonic-clonic, abnormal movement) – brain edema – increased intracranial tension (ICT) – CNS infection.



NB Cow milk allergy: Ag - Ab reaction within 2-3 weeks
skin changes. Atrophic changes of villi - bloody stool -
bad general condition. - chest problems.

Management in Out-patient Department:

[1] Diarrhea with no dehydration:

Counsel the mother on 3 rules for home treatment:

1. Give extra fluids:

- Teach the mother how to mix and give oral rehydration solution (ORS)/200 mL water. *Hydrosafe - Lohydram - Rehydrosine*
- Slowly by cup and spoon every 1-2 min or by bottle (slowly).
- Up to 2 years 50-100 mL after each loose/watery motion.
- Two years or more 100-200 mL after each loose watery stool.

2. Continue feeding:

- If breast feeding (BF) continue.
- If not exclusive BF, give food based fluids; soup, Rice water or artificial milk if mother can afford.

- Sugar free diet.

3. When to return back for consultation:

- Drinks more than 4-5 ORS packets/day.
- Drinks poorly or unable to drink.
- High fever - rapid breathing.
- Bloody stool.
- Severe abdominal distention.

[2] Diarrhea with some dehydration:

Refer to the oral section to receive ORS for 2-4 hours.

- About 50-80 mL/kg over 2-4 hours by cup and spoon/every 1-2 min or frequent sips from cup for older children.
- Check regularly to see improvement or any problems.
- If child vomits, give ORS very slowly.
- If child's eye lids become puffy, stop ORS.
- Reassess within 4 hours, still has some dehydration add more ORS.
- If mother has to leave before 4 hours, teach her to complete rehydration at home. Instruct her 3 rules, and when to return.
- Advice breast feeding if the child wants.

If gets worse shift to IV fluid (failure of ORS):

- Reassess ... within 4-6 hours and classify the dehydration again then treat accordingly:

post. FT
x - Torsk
x + femoral - Inadequacy.

eyes
Tongue skin
extremities
Breathing
Normal / cold / cyanotic
shocked.
Take a fold of sc fat.

a. No Dehydration:

Discharge

- If possible, observe the child for few hours to make sure that mother can maintain hydration by giving ORS.
- Discharge from out-patient clinic and oral rehydration section by 3 golden rules.

b. Sill dehydration

- If able to drink manage as some dehydration till discharge.
- If failed ORS, shift to I.V fluids/consider admission.

c. Complication

- Deteriorating general condition.
- Altered mental status.
- Coffee ground vomitus.
- Convulsion.
- Severe abdominal distention, intestinal obstruction?
- Bleeding tendency.
- Severe respiratory distress.
- Electrolyte disturbance
 - * Severe hypernatremia, $\text{Na} > 160 \text{ mEq/L}$ or hyponatremia $< 120 \text{ mEq/L}$.
 - * Severe hypokalemia $< 2 \text{ mEq/L}$.
 - * Severe hypoglycemia or hyperglycemia (usually occurs in severe dehydration/shock).
 - * Severe metabolic acidosis (occurs in severe condition).

Admit to in-patient section

[3] Severe dehydration ± shock:

- Start IV fluid immediately.
- If possible drain blood sample before set up the line for serum Na^+ , K^+ , creat, BUN, glucose, blood gases.
- If difficulty to set up IV, insert nasogastric tube with ORS, until IV line is in place or use intraosseous cannulation.
- Solutions used for IV fluid.

Polyvalent, Pansol, Methahydrane (100 mL/kg) divided as follows:

Age	First give 30 mL/kg in	Then give 70 mL/kg in
Infants under 12 m	1 hour	5 hours
Children (12 m – up to 5 years)	30 min	2.5 hours

- If the child is able to drink treat accordingly, you may try some sips of ORS or breast milk.
- If the pulse is very weak or not detected after antishock therapy (**consider admission to ICU**).

Management at In-patient Department

Cases for admission:

- Shocked cases not responding to management in OP clinic after antishock measures (30 mL/kg/half-one hour).
- Complications associated with diarrhea.
 1. Diarrhea + pneumonia.
 2. Sepsis with high fever, bleeding tendency.
 3. Bloody diarrhea with bad general condition.
 4. Severe electrolytes disturbance.
 5. Severe abdominal distention.
 6. Complicated persistent diarrhea.
 7. Bleeding tendency, coffee ground vomitus, ecchymosis.
 8. Suspected intracranial tension (ICT), neurological disorders e.g., convulsions.

Work up and investigation at in-patients dept:

- Admission sheets, follow up sheets and progress notes should be filled up.
- Routine investigations for all admitted cases:
 1. Stool analysis routinely, stool culture/sensitivity, if needed.
 2. Serum electrolytes "Na⁺, K⁺, Ca⁺⁺" and glucose.
 3. Blood gases.
 4. Renal functions (BUN – Creat).
 5. CBC – CRP.
 6. Others, if indicated (liver functions, PT, PC, T. protein, Albumin, ...)

Treatment of Complications:

* Hypokalaemia:

(Serum K⁺ < 2.5-3 mEq/L)

- **When to suspect:**
 1. Severe losses (mainly vomiting).
 2. Abdominal distention.
 3. Severe losses + protein energy malnutrition (PEM).
- If serum K⁺ ≤ (2.5-3 mEq/L) and child is asymptomatic, give oral potassium, 1-4 mEq/kg/24 hr
- If serum K⁺ < (2-2.5 mEq/L) → I.V rout.
K⁺ required use this equation:

$$\frac{\text{B. wt.} \times \text{deficit (3.5-serum K}^+ \text{ level)} \times 0.6}{2} = \text{K needed for replacement}$$

Add it to I.V therapy, (Never direct I.V).

- **OR** : use I.V solution containing K 40 mEq/L, or 60 mEq/L in severe hypokalaemia.

MONITOR Serum K closely.



normal saline 0.9% (1 meq \rightarrow 6.6 cm)
hypertonic saline 2.4% (1 meq \rightarrow 2.2 cm)

* Hyponatremia:

- Serum Na (<120-125 mEq/L).
- **When to suspect:**
 1. Severe dehydration.
 2. Shock.
 3. Severe losses + PEM.
 4. Disturbed conscious level.
- If (<120 mEq/L) give hypertonic solution (Saline 2.7%) to raise serum Na as follows:

$\frac{1}{2}$ Normal sol. (1 meq \rightarrow 6.6 cm x 2)
0.45

$(125 - \text{serum Na}) \times 0.6 = \text{total mEq, needed to correct serum Na (over 2-3 hours)}$.

Then if serum Na become 120 mEq/L or more manage accordingly.

* Hypernatremic dehydration:

- Serum Na (>150 mEq/L).
- **When to suspect:**
 1. History of faulty mixing ORS, or increase intake prior to attending the hospital >5 packets/day.
 2. Severe dehydration.
 3. Irritability – altered mental status, convulsions.
- If serum Na⁺ (150-160 mEq/L) ORS is effective. Such cases can be treated in out-patient unless accompanied with other complications.
- If Na⁺ 160 mEq/L or more (solution used not more than 60 mEq/L) i.e., Kadlex: Na HCO₃ (8.4%)
15 : 1
470 cc : 30 cc
In a rate of 100-130 mL/kg/ 24 hours, according to dehydration, status.
- Monitor serum Na every (8-12 hours).
- If serum Na decreases by 0.5-1 mEq/L \approx 12 mEq/ 24 hours, you are on proper path.
- Sips of plain water can be given orally by 5 mL/kg/day during IV therapy.

Reassess:

- If Na becomes within normal, manage according to clinical evaluation.
- Still high (>160 mEq/L) continue IV fluids with Na not more than 60 mEq/L.
- Up to 160 mEq/L, no dehydration, stable, ... tolerate oral, shift to oral rehydration .. with ORS for ongoing losses and start feeding.
- If child developed convulsions, or renal impairment:

Consult nephrologists.

Consult neurologist.

- Rule out; CNS infection or ↑ICT.
- If serum Na still high consider peritoneal dialysis.

*** Metabolic acidosis:**

- The serum HCO_3 and PH fall and PCO_2 decreases.
- Metabolic acidosis due to diarrhea and dehydration is usually with normal anion gap.
 $\text{Anion gap} = (\text{Na} + \text{K}) - (\text{CL} + \text{HCO}_3)$.
- Normal anion gap = 12 mEq/L with range 8-16 mEq/L
- If $\text{HCO}_3 < 5$ mEq/L give Na HCO_3 (8.4%).
 $(12 - \text{serum } \text{HCO}_3) \times \text{BW} \times 0.3 = (\text{HCO}_3 \text{ deficit})$
- If HCO_3 10-15 mEq/L and PH less than 7.2 give Na HCO_3 . if PH > 7.2 correct dehydration and reassess.

*** Convulsion:**

- Initially correct metabolic causes (hyper Na^+ , hypo Na^+ , hypo Ca^{++} , hypoglycemia).
- If recurrent, diazepam IV or rectally by 0.1 mg/kg, or Phenobarbital loading dose 20 mg/kg IV slowly.
- If recurrent think of Encephalopathy, CNS infection.
- Consult seniors for, 1-CSF 2-CT brain 3-EEG.

*** Hyperglycaemia:**

- Bt glucose may reach > 400 mg/dL or more, it is considered as (stress hyperglycemia), due to release of stress hormones, that block insulin action.
- After correction of dehydration and shock, if hyperglycemia persists, **consult diabetologist for insulin therapy.**

*** Respiratory distress:**

- May be due to:
 1. Shock.
 2. Intra-pulmonary Hg.
 3. Severe infection
 4. Septicemia.
 5. Metabolic acidosis.

Treat accordingly

*** Bleeding:**

I. Hypoprothrombenemia:

- Bleeding from venipuncture.
- PT, PTT, high
- Plat count, ----- low.
Give vit K 1-2 mg/IM

II. DIC:

- The previous finding +
- Severe anemia.
- ± coffee ground vomiting.
- Purpuric eruption.
- FDP +++ (Fibrin degradation products).

Fresh blood transfusion/fresh frozen plasma, aggressive antibiotics, maintain hydration, treat other complications.

III. HUS:

- Bleeding tendency.
- Severe anemia.
- ± Hematuria.
- ± impaired renal function
- Liver function ++
- Hepatosplenomegaly.

Consult Nephrologist:

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Feeding after rehydration

- If child is on breast milk, continue as the child able to drink.
- If child on artificial milk $\frac{1}{2}$ conc. – then increase gradually to full concentration.
- If child 6 m or older; soft foods (soup, mashed chicken, vegetables yoghurt, cereals,).
- If diarrhea continues with feeding and milk, suspect lactose intolerance: Clinitest +ve for disaccharides or reducing substances in stool → Give lactose free formula.
- If diarrhea persist consider elemental formula or partial parenteral nutrition.

Drugs and diarrhea

- Anti-diarrheal drugs → ^{as ↑ incidence of persistent diarrhea.} no role in management of diarrhea. *e.g. Antinal, smecta,*
- Anti-emetics → limited role in treating persistent vomiting.

Antimicrobial and diarrhea:

Should not be given routinely to children with diarrhea.

Only in:

- Bloody diarrhea with bad general condition.
- Suspected cases of Gardia, Amoeba.
- Diarrhea associated with extra-intestinal infection, *e.g., pharyngitis, otitis media, skin infection, chest infection, septicemia*
3rd generation cephalosporin or amoxicillin , And/or Metronidazole,
..... are the drugs of choice.

N.B.: Usually start by single antibiotic.

Clinical particulars (in out-patient clinic):

[I] Assessment of diarrhea patients for dehydration:

	No dehydration	Some dehydration	Severe dehydration
Look at: The case	Well, alert.	Restless, irritable	Lethargic, unconsciousness.
Eyes	Normal.	Sunken eyes.	Sunken eyes.
Thirst	Drinks normally, not thirsty.	Thirsty, drinks eagerly.	Drinks poorly, or unable to drink.
Feel: Skin pinch	Goes back quickly	Goes back slowly	Goes back very slowly
Decide	Patients has no dehydration	Has two or more signs, there is some dehydration	Has two or more signs of the above, it is severe dehydration
Treat	Give three golden rules: - Extra fluids /ORS - Continue feeding - When to return?	Weight the patient and treat at OP section with ORS by cup and spoon.	Weigh the patient and start IV fluids.

[II] Investigations can be carried out in OP clinic:

- Stool analysis only for bloody diarrhea and persistent diarrhea >14 days.
- Serum electrolytes: Na – K⁺, Ca⁺⁺ and glucose only for severe dehydration & shock and dehydration with convulsion.
- Kidney functions BUN, Creat for severe dehydration shock, decreased urinary flow.
- CBC only for pallor, bad general condition and fever.

[III] For discharged cases after rehydration:

Consider 3 rules + medications required according to clinical conditions, e.g.:

- * Antipyretics → paracetamol (10-15 mg/kg/dose). *no supp.*
- * Antiparasitic → metronitazole (15-30 mg/kg/dose) *Amebiac 10 days 18 or 16 hrs*
→ nitazoxanide 7.5 mg/kg/dose *Giardia 7 days 12 hrs*
- Antibiotics should not be given as a routine for acute diarrhea.
- Zinc supplementation (1-3mg/kg) for 10-14 days with other micronutrients.
- Pre biotics and pro biotics can help in shortening diarrhea duration.
- Contact senior unit members for any enquiry.
- Amoxicillin
- 3rd generation cephalosporin } Drug of choice, if indicated
- Azithromycin for (shigellosis & campylobacter).

* if fever, offensive, bloody stool, pus cells, RBCs → Bacterial

(Sulphamethoxazole + Trimethoprim) septazole or Sutrim (6 Mo - 1

>1 15/100 for 7 Days

* if viral *Anti Rotavirus or Meta zoxide: NB used in using 6*

Antigiardiasis but e viral ↓ duration, ↓ Freq. *Interferon.*

7.5 mg/kg/dose (100mg/5ml) for 3 days

* Vomiting (due to dehydration & feeding) *primarily extra Δ, not dose related*

regulation of feeding. *2ml x 5.*
Motinorm → regulate intest. mov., Cortigen

*low grade fever
↑ freq.
No offensive
No bloody
Vomiting*

References:

1. Koletzko S, Osterrieder S. Acute infectious diarrhea in children. *Dtsch Arztebl Int* 2009 Aug; 106 (33): 539-547; quiz 548 Epub 2009 Aug 14.
2. Richards L, Claeson M, Pierce NF. Management of acute diarrhea in children : lessons learned. *Pediatr Infect Dis J*. 1993 Jan; 12 (1): 5-9.
3. Santosham M, Fayad I, Abu Zikri M, Hussein A, Amponsah A, Duggan C, Hashem M, El-Sady N, Abu Zikri M, Fontaine O. A double-blind clinical trial comparing world health organization oral rehydration solution with a reduced osmolarity solution containing equal amounts of sodium and glucose. *J Pediatr* 1996 Jan.; 128 (1): 45-51.
4. Rapid intravenous rehydration in acute diarrhea. Sperotto G. *Bol Med Hosp Infant Mex* 1992 Aug; 48 (8): 506-513.
5. The clinical epidemiology of hypernatremia in diarrhea during treatment with oral rehydration in Egypt. Ahmed Youssef, IM Fayyad, GJ Ebrahim. *Trop Ped* Dec 1988; 34: 289-293.
6. Diarrheal disease morbidity and home treatment practices in Egypt. Jousilahti P, Madkour SM, Lambrechts T, Sherwin E. *Public Health* 1997 Jan; 111 (1): 5-10.
7. Oral rehydration therapy for diarrhea in children – a basic primer. Hirschitov N. *Nutr Rev* 1982 Apr; 40 (4): 97-104. Review.
8. Rational home management of diarrhea. Almroth S, Latham MC. *Lancet* 1995 Mar 18; 345 (8951): 709-711.
9. Oral rehydration therapy for diarrhea: an example of reverse transfer of technology. Santosham M, Keenan EM, Tulloch J, Broun D, Glass R. *Pediatrics* 1997 Nov; 100 (5): E10.
10. Rational management of diarrhea in children. World Health Organization (WHO). Division of diarrheal and acute respiratory disease control. *Essent Drugs Monit* 1991; 11: 10-11.
11. Acute diarrheal diseases – an approach to management. Sur D, Bhattacharya SK. *J Indian Med Assoc* 2006 May; 104 (5) 220-223.
12. The management of acute diarrhea in children: oral rehydration, maintenance, and nutritional therapy. Centers for Disease Control and Prevention. Duggan C, Santosham M, Glass RI. *MWWWR Recomm Rep* 1992 Oct 16; 41 (RR-16): 1-20.
13. Oral rehydration therapy for invasive diarrhea. Varavithya W, Sunthomkachit R, Eampokalap B. *Rev Infect Dis* 1991 Mar-Apr; 13 (Suppl.4): S325-S331. Review.

14. Trends in the management of childhood diarrhea in Egypt 1979-1990. Miller PC. J diarrheal Dis Res 1992 Dec; 10 (4): 193-200.
15. Feeding infants and young children with acute diarrhea. Chouraqui JP, Michard-Lenoir AP. Arch Pediatr 2007 Oct ; 14 (Suppl.3) : S176-S180.
16. Substantial reduction in severe diarrheal morbidity by daily zinc supplementation in young north Indian children. Bhandari N, Bahl R, Taneja S, Strand T, Molbak K, Ulvik RJ, Sommerfeitt H, Bhan MK. Pediatrics 2002 Jun; 109 (6): E66.
17. Dealing with diarrhea. Fontaine O. Child Health Dialogue 1996 (3-4): 5.
18. Oral rehydration solutions and acute diarrhea: an update. Mouterde O. Arch Pediatr 2007 Oct; 14 (Suppl.3): S165-S168.
19. Oral rehydration therapy of infantile diarrhea: a controlled study of well-nourished children hospitalized in the United States and Panama. Santosham M, Daum RS, Dillman L, Rodriguez JL, Luque S, Russell R, Kourany M, Ryder RW, Bantlett AV, Rosenberg A, Benenson AS, Sack RB. N Engl J Med 1982 May 6; 306 (18): 1070-1076.
20. The effect of oral rehydration solution and recommended home fluids on diarrhea mortality. Muros MK, Waker CL, Black RE. Int J Epidemiol 2010 Apr; 39 (Suppl.1): 75-87.
21. Zinc with oral rehydration therapy reduces stool output and duration of diarrhea in hospitalized children: a randomized controlled trial. Bhatnagar S, Bahl R, Sharma PK, Kumar GT, Saxena SK, Bhan MK. J Pediatr Gastroenterol Nutr 2004 Jan; 38 (1): 34-40.
22. The treatment of diarrhea. A manual for physicians. World Health Organization 2005.
23. Joint Statement Clinical Management of Acute Diarrhea. These new recommendations, formulated by UNICEF and WHO in collaboration with USAID and experts worldwide, take into account new research findings while building on past recommendations. UNICEF/WHO (2004).

ما هي البريبايوتكس ؟

البريبايوتكس هي ألياف تعمل على تحفيز نمو البكتيريا النافعة في الأمعاء. تحتوي أمعاء الطفل بشكل طبيعي على بكتيريا نافعة وبكتيريا ضارة.

لماذا يحتاج الطفل إلى البكتيريا النافعة في أمعائه ؟

يحتاج الطفل إلى البكتيريا النافعة وذلك لأنها:

- ★ تحفز جهاز المناعة .
 - ★ تمنع نمو البكتيريا الضارة .
 - ★ تساعد في علاج الحساسية.
 - ★ تساعد في الهضم حيث تعمل على جعل البراز أكثر طراوة.
 - ★ تساعد على العلاج والوقاية من حالات الإسهال للأطفال.
- بتحفيز البكتيريا النافعة لدى الطفل عن طريق البريبايوتكس تضمن حصوله على الفوائد التي تقدمها

What are Prebiotics?

Prebiotics are fibres which stimulate the growth of the friendly bacteria in the intestine.

The baby and toddler's intestines naturally contain friendly bacteria and unfriendly bacteria.

Why is the friendly bacteria needed in the intestins?

the friendly bacteria is needed because :

1. it stimulates the immune system.
2. it prevents the growth of unfriendly bacteria.
3. it helps in the treatment and prevention of diarrhea.

So by stimulating the bebys good bacteria with prebiotics you ensure that its benefits are delivered to his / her body.

بيبيلاك Bebelac®

**بيبيلاك
Bebelac®**

**بيبيلاك بريماتيور
Premator Bebelac**

**بيبيلاك ١
Bebelac®**

**منذ الولادة وما بعدها
From birth onwards**

**بيبيلاك إي سي
Bebelac®**

**وخاصة في حالات المغص
والتقلصات للرضع
Especially in cases of colic
& contractions of infants**

**بيبيلاك ٢
Bebelac®**

**من سن ٦ أشهر
From 6 months**

**بيبيلاك إف إل
Bebelac®**

**في حالات الإسهال
الحاد والمزمن
In Cases of diarrhea
Acute & chronic**

**بيبيلاك ٣
Bebelac®**

**من ١ إلى ٤ سنوات
From 1 To 4 Years**